

Isothermal (vapor + liquid) equilibria and excess enthalpy data of {1-hexene + methyl butyl ether (MBE)} and {1-hexene + methyl tert-butyl ether (MTBE)} binary systems at several temperatures

- [Rachida Hani^a](#),
 - [Roland Solimando^b](#),
 - [Latifa Negadi^c](#),   
 - [Jacques Jose^d](#),
 - [Ahmed Ait Kaci^a](#)
- ^a LTMM, Laboratoire de Thermodynamique et Modélisation Moléculaire, Université des Sciences et Technologie Houari Boumediene, P.O. Box 132, 16111 El Alia, Algeria
 - ^b Laboratoire de Thermodynamique des Milieux Polyphasés, ENSIC – INPL, 1 rue Grandville BP 451, 54001 Nancy Cedex, France
 - ^c LATA2M, Laboratoire de Thermodynamique Appliquée et Modélisation Moléculaire, University AbouBekr Belkaid of Tlemcen, P.O. Box 119, Tlemcen 13000, Algeria
 - ^d LSA, Laboratoire des Sciences Analytiques, CNRS-UMR 5280, Université Claude Bernard, Lyon I. 43, Bd du 11 Novembre 1918, Villeurbanne Cedex 69622, France

Abstract

The vapor pressures of {1-hexene + methyl butyl ether (MBE)} and {1-hexene + methyl tert-butyl ether (MTBE)} binary mixtures and of the three pure components were measured by means of a static device at temperatures between (263 and 333) K. The data were correlated with the Antoine equation. From these data, excess Gibbs functions were calculated for several constant temperatures and fitted to a third-order Redlich–Kister equation using the Barker's method. Additionally, molar excess enthalpies, H^E , for the two binary systems have been measured at 303.15 K using an isothermal flow calorimeter.

Keywords

- (Vapor + liquid) equilibria;
- 1-Hexene;
- Ether;
- Excess Gibbs free energy;
- Excess enthalpy